arXiv:1111.3115v2 [math.DG] 28 Dec 2011

Estimates for Eigenvalues of Poly-harmonic Operators

Guoxin Wei and Lingzhong Zeng

Abstract

In this paper, we study eigenvalues of the poly-Laplacian with arbitrary order on a bounded domain in an n-dimensional Euclidean space and obtain a lower bound for eigenvalues, which generalizes the results due to Cheng-Wei [5] and gives an improvement of results due to Cheng-Qi-Wei [3].

Keywords: the eigenvalue problem; a lower bound for eigenvalues; the ploy-Laplacian with arbitrary order.

2010 Mathematics Subject Classification: 35P15.

1 Introduction

Let Ω be a bounded domain with piecewise smooth boundary $\partial\Omega$ in an n-dimensional Euclidean space $\mathbb{R}^n$. Let λ_i be the i-th eigenvalue of Dirichlet eigenvalue problem of the poly-Laplacian with arbitrary order:

$$\begin{cases}(-\Delta)^l u = \lambda u, & in\ \Omega,\\ u = \dfrac{\partial u}{\partial \nu} = \cdots = \dfrac{\partial^{l-1} u}{\partial \nu^{l-1}} = 0, & on\ \partial\Omega,\end{cases} \tag{1.1}$$

where Δ is the Laplacian in $\mathbb{R}^n$ and ν denotes the outward unit normal vector field of the boundary $\partial\Omega$. It is well known that the spectrum of this eigenvalue problem (1.1) is real and discrete:

$$0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \to +\infty,$$

where each λ_i has finite multiplicity which is repeated according to its multiplicity. Let $V(\Omega)$ denote the volume of Ω and let B_n denote the volume of the unit ball in $\mathbb{R}^n$.

When $l = 1$, the eigenvalue problem (1.1) is called a fixed membrane problem. In this case, one has the following Weyl's asymptotic formula

$$\lambda_k \sim \frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad k \to +\infty. \tag{1.2}$$

From the above asymptotic formula, one can derive

$$\frac{1}{k}\sum_{i=1}^{k} \lambda_i \sim \frac{n}{n+2} \frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad k \to +\infty. \tag{1.3}$$

Pólya [12] proved that

$$\lambda_k \geq \frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad \text{for } k = 1, 2, \cdots, \tag{1.4}$$

if Ω is a tiling domain in $\mathbb{R}^n$. Furthermore, he proposed a conjecture as follows:

Conjecture of Pólya. *If* Ω *is a bounded domain in* $\mathbb{R}^n$*, then the* k*-th eigenvalue* λ_k *of the fixed membrane problem satisfies*

$$\lambda_k \geq \frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad \text{for } k = 1, 2, \cdots . \tag{1.5}$$

On the conjecture of Pólya, Berezin [2] and Lieb [9] gave a partial solution. In particular, Li and Yau [8] proved that

$$\frac{1}{k}\sum_{i=1}^{k} \lambda_i \geq \frac{n}{n+2}\frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad \text{for } k = 1, 2, \cdots . \tag{1.6}$$

The formula (1.3) shows that the result of Li and Yau is sharp in the sense of average. From this formula (1.6), one can infer

$$\lambda_k \geq \frac{n}{n+2}\frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}}, \quad \text{for } k = 1, 2, \cdots , \tag{1.7}$$

which gives a partial solution for the conjecture of Pólya with a factor $\dfrac{n}{n+2}$. Recently, Melas [10] has improved the estimate (1.6) to the following:

$$\frac{1}{k}\sum_{i=1}^{k} \lambda_i \geq \frac{n}{n+2}\frac{4\pi^2}{(B_n V(\Omega))^{\frac{2}{n}}} k^{\frac{2}{n}} + \frac{1}{24(n+2)}\frac{V(\Omega)}{I(\Omega)}, \quad \text{for } k = 1, 2, \cdots , \tag{1.8}$$

where

$$I(\Omega) = \min_{a\in\mathbb{R}^n} \int_\Omega |x-a|^2 dx$$

is called *the moment of inertia* of Ω.

When $l = 2$, the eigenvalue problem (1.1) is called a clamped plate problem. For the eigenvalues of the clamped plate problem, Agmon [1] and Pleijel [11] obtained

$$\lambda_k \sim \frac{16\pi^4}{(B_n V(\Omega))^{\frac{4}{n}}} k^{\frac{4}{n}}, \quad k \to +\infty. \tag{1.9}$$

From the above formula (1.9), one can obtain

$$\frac{1}{k}\sum_{i=1}^{k} \lambda_i \sim \frac{n}{n+4}\frac{16\pi^4}{(B_n V(\Omega))^{\frac{4}{n}}} k^{\frac{4}{n}}, \quad k \to +\infty. \tag{1.10}$$

Furthermore, Levine and Protter [7] proved that the eigenvalues of the clamped plate problem satisfy the following inequality:

$$\frac{1}{k}\sum_{i=1}^{k} \lambda_i \geq \frac{n}{n+4}\frac{16\pi^4}{(B_n V(\Omega))^{\frac{4}{n}}} k^{\frac{4}{n}}. \tag{1.11}$$

The formula (1.10) shows that the coefficient of $k^{\frac{4}{n}}$ is the best possible constant. By adding to its right hand side two terms of lower order in k, Cheng and Wei [4] obtained the following estimate which is an improvement of (1.11):

$$\begin{aligned}\frac{1}{k}\sum_{i=1}^{k}\lambda_i \geq \frac{n}{n+4}\frac{16\pi^4}{(B_nV(\Omega))^{\frac{4}{n}}}k^{\frac{4}{n}} &\\ +\left(\frac{n+2}{12n(n+4)}-\frac{1}{1152n^2(n+4)}\right)\frac{4\pi^2}{(B_nV(\Omega))^{\frac{2}{n}}}\frac{n}{n+2}\frac{V(\Omega)}{I(\Omega)}k^{\frac{2}{n}} &\\ +\left(\frac{1}{576n(n+4)}-\frac{1}{27648n^2(n+2)(n+4)}\right)\left(\frac{V(\Omega)}{I(\Omega)}\right)^2.&\end{aligned} \tag{1.12}$$

Very recently, Cheng and Wei [5] have improved the estimate (1.12) to the following:

$$\begin{aligned}\frac{1}{k}\sum_{i=1}^{k}\lambda_i \geq \frac{n}{n+4}\frac{16\pi^4}{(B_nV(\Omega))^{\frac{4}{n}}}k^{\frac{4}{n}} &\\ +\frac{n+2}{12n(n+4)}\frac{4\pi^2}{(B_nV(\Omega))^{\frac{2}{n}}}\frac{n}{n+2}\frac{V(\Omega)}{I(\Omega)}k^{\frac{2}{n}} &\\ +\frac{(n+2)^2}{1152n(n+4)^2}\left(\frac{V(\Omega)}{I(\Omega)}\right)^2.&\end{aligned} \tag{1.13}$$

When l is arbitrary, Levine and Protter [7] proved the following

$$\frac{1}{k}\sum_{i=1}^{k}\lambda_i \geq \frac{n}{n+2l}\frac{\pi^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}}, \quad \text{for} \quad k=1,2,\cdots, \tag{1.14}$$

which implies that

$$\lambda_k \geq \frac{n}{n+2l}\frac{\pi^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}}, \quad \text{for} \quad k=1,2,\cdots. \tag{1.15}$$

By adding l terms of lower order of $k^{\frac{2l}{n}}$ to its right hand side, Cheng, Qi and Wei [3] obtained more sharper result than (1.14):

$$\begin{aligned}\frac{1}{k}\sum_{i=1}^{k}\lambda_i \geq \frac{n}{n+2l}\frac{(2\pi)^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}} + \frac{n}{(n+2l)} &\\ \times\sum_{p=1}^{l}\frac{l+1-p}{(24)^p n\cdots(n+2p-2)}\frac{(2\pi)^{2(l-p)}}{(B_nV(\Omega))^{\frac{2(l-p)}{n}}}\left(\frac{V(\Omega)}{I(\Omega)}\right)^p k^{\frac{2(l-p)}{n}}.&\end{aligned} \tag{1.16}$$

In this paper, we investigate eigenvalues of the Dirichlet eigenvalue problem (1.1) of Laplacian with arbitrary order and prove the following:

Theorem 1.1. *Let Ω be a bounded domain in an n-dimensional Euclidean space $\mathbb{R}^n$. Assume that $l \geq 2$ and λ_i is the i-th eigenvalue of the eigenvalue problem (1.1). Then the eigenvalues satisfy*

$$\begin{aligned}\frac{1}{k}\sum_{j=1}^{k}\lambda_j \geq \frac{n}{n+2l}\frac{(2\pi)^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}} &\\ +\frac{l}{24(n+2l)}\frac{(2\pi)^{2(l-1)}}{(B_nV(\Omega))^{\frac{2(l-1)}{n}}}\frac{V(\Omega)}{I(\Omega)}k^{\frac{2(l-1)}{n}} &\\ +\frac{l(n+2(l-1))^2}{2304n(n+2l)^2}\frac{(2\pi)^{2(l-2)}}{(B_nV(\Omega))^{\frac{2(l-2)}{n}}}\left(\frac{V(\Omega)}{I(\Omega)}\right)^2 k^{\frac{2(l-2)}{n}}.&\end{aligned} \tag{1.17}$$

Remark 1.1. *When $l = 2$, Theorem* 1.1 *reduces to the result of Cheng-Wei* [5].

Remark 1.2. *When $l \geq 2$, we give an important improvement of the result* (1.16) *due to Cheng-Qi-Wei* [3] *since the inequality* (1.17) *is sharper than the inequality* (1.16). *About this fact, we will give a proof in Section* 3.

2 A Key Lemma

In this section, we will give a key Lemma which will play an important role in the proof of Theorem 1.1.

Lemma 2.1. *Let $b \geq 2$ be a positive real number and $\mu > 0$. If $\psi : [0,\ +\infty) \to [0,\ +\infty)$ is a decreasing function such that*

$$-\mu \leq \psi'(s) \leq 0$$

and

$$A := \int_0^\infty s^{b-1}\psi(s)ds > 0,$$

then, for any positive integer $l \geq 2$, we have

$$\begin{aligned}\int_0^\infty s^{b+2l-1}\psi(s)ds \geq\ & \frac{1}{b+2l}(bA)^{\frac{b+2l}{b}}\psi(0)^{-\frac{2l}{b}} \\ & + \frac{l}{6b(b+2l)\mu^2}(bA)^{\frac{b+2(l-1)}{b}}\psi(0)^{\frac{2b-2l+2}{b}} \\ & + \frac{l(b+2(l-1))^2}{144b^2(b+2l)^2\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}.\end{aligned} \tag{2.1}$$

Proof. Let

$$\varrho(t) = \frac{\psi\big(\frac{\psi(0)}{\mu}t\big)}{\psi(0)}, \tag{2.2}$$

then we have $\varrho(0) = 1$ and $-1 \leq \varrho'(t) \leq 0$. Without loss of generality, we can assume

$$\psi(0) = 1 \text{ and } \mu = 1.$$

Define

$$D_l := \int_0^\infty s^{b+2l-1}\psi(s)ds.$$

One can assume that $D_l < \infty$, otherwise there is nothing to prove. Since $D_l < \infty$, we can conclude that

$$\lim_{s\to\infty} s^{b+2l-1}\psi(s) = 0.$$

Putting $h(s) = -\psi'(s)$ for $s \geq 0$, we get

$$0 \leq h(s) \leq 1 \ \text{ and } \ \int_0^\infty h(s)ds = \psi(0) = 1.$$

By making use of integration by parts, one has

$$\int_0^\infty s^b h(s)ds = b\int_0^\infty s^{b-1}\psi(s)ds = bA,$$

and

$$\int_0^\infty s^{b+2l}h(s)ds \leq (b+2l)D_l,$$

since $\psi(s) > 0$. By the same assertion as in [10], one can infer that there exists an $\epsilon \geq 0$ such that

$$\int_{\epsilon}^{\epsilon+1} s^b ds = \int_0^{\infty} s^b h(s) ds = bA, \tag{2.3}$$

and

$$\int_{\epsilon}^{\epsilon+1} s^{b+2l} ds \leq \int_0^{\infty} s^{b+2l} h(s) ds \leq (b+2l) D_l. \tag{2.4}$$

Let

$$\Theta(s) = bs^{b+2l} - (b+2l)\tau^{2l}s^b + 2l\tau^{b+2l} - 2l\tau^{b+2(l-1)}(s-\tau)^2,$$

then we can prove that $\Theta(s) \geq 0$. By integrating the function $\Theta(s)$ from ϵ to $\epsilon+1$, we deduce from (2.3) and (2.4), for any $\tau > 0$,

$$b(b+2l)D_l - (b+2l)\tau^{2l}bA + 2l\tau^{b+2l} \geq \frac{l}{6}\tau^{b+2(l-1)}. \tag{2.5}$$

Define

$$f(\tau) := (b+2l)\tau^{2l}bA - 2l\tau^{b+2l} + \frac{l}{6}\tau^{b+2(l-1)},$$

then we can obtain from (2.5) that, for any $\tau > 0$,

$$D_l = \int_0^{\infty} s^{b+2l-1}\psi(s) ds \geq \frac{f(\tau)}{b(b+2l)}.$$

Taking

$$\tau = (bA)^{\frac{1}{b}} \left(1 + \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{1}{b}},$$

then one has

$$\begin{aligned} f(\tau) = (bA)^{\frac{b+2l}{b}} &\left(b - \frac{l(b+2(l-1))}{6(b+2l)}(bA)^{-\frac{2}{b}}\right)\left(1 + \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{2l}{b}} \\ &+ \frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}\left(1 + \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{b+2(l-1)}{b}}. \end{aligned} \tag{2.6}$$

Next, we consider four cases:

Case 1: $b \geq 2l$. For $t > 0$, we have from the Taylor formula

$$\begin{aligned} (1+t)^{\frac{2l}{b}} \geq 1 + \frac{2l}{b}t + \frac{2l(2l-b)}{2b^2}t^2 + \frac{2l(2l-b)(2l-2b)}{6b^3}t^3 \\ + \frac{2l(2l-b)(2l-2b)(2l-3b)}{24b^4}t^4 \end{aligned}$$

and

$$\begin{aligned} (1+t)^{\frac{b+2(l-1)}{b}} \geq 1 + \frac{2(l-1)+b}{b}t + \frac{(2(l-1)+b)(l-1)}{b^2}t^2 \\ + \frac{(2(l-1)-b)(l-1)(2(l-1)+b)}{3b^3}t^3. \end{aligned}$$

Putting $t = \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}$, we have from $(bA)^{\frac{2}{b}} \geq \frac{1}{(b+1)^{\frac{2}{b}}} \geq \frac{1}{3} > \frac{1}{4}$ (also see [5]) that $t < \frac{1}{3}$ and $b - 2lt > \frac{4l}{3} > 0$. And then, we obtain

$$\begin{aligned} &\left(b - \frac{l(b+2(l-1))}{6(b+2l)}(bA)^{-\frac{2}{b}}\right)\left(1 + \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{2l}{b}} \\ &= (b-2lt)(1+t)^{\frac{2l}{b}} \end{aligned}$$

$$\begin{aligned}
&\geq (b-2lt)\left[1+\frac{2l}{b}t+\frac{2l(2l-b)}{2b^2}t^2+\frac{2l(2l-b)(2l-2b)}{6b^3}t^3\right.\\
&\quad\left.+\frac{2l(2l-b)(2l-2b)(2l-3b)}{24b^4}t^4\right]\\
&\geq b-\frac{l(2l+b)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\\
&\quad-\frac{(2l-b)(8l^2+4lb)}{6b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\quad-\frac{(2l-b)(2l-2b)(12l^2+6lb)}{24b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4
\end{aligned}$$

and

$$\begin{aligned}
&\left(1+\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{b+2(l-1)}{b}}\\
&=(1+t)^{\frac{b+2(l-1)}{b}}\\
&\geq 1+\frac{2(l-1)+b}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)\\
&\quad+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\\
&\quad+\frac{(2(l-1)-b)(l-1)(2(l-1)+b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
f(\tau)&=(b+2l)\tau^{2l}bA-2l\tau^{b+2l}+\frac{l}{6}\tau^{b+2(l-1)}\\
&\geq (bA)^{\frac{b+2l}{b}}\left[b-\frac{l(2l+b)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\right.\\
&\quad-\frac{(2l-b)(8l^2+4lb)}{6b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\quad\left.-\frac{(2l-b)(2l-2b)(12l^2+6lb)}{24b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4\right]\\
&\quad+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}\left[1+\frac{2(l-1)+b}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)\right.\\
&\quad+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\\
&\quad\left.+\frac{(2(l-1)-b)(l-1)(2(l-1)+b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\right]\\
&=b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}+\eta_1,
\end{aligned}$$

where

$$\begin{aligned}\eta_1 = {} & \frac{2l(l+b-3)(b+2l)}{3b^2}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-6}{b}} \\ & + \frac{l(b+2(l-1))(4(l-1)(2(l-1)-b)-3(2l-b)(l-b))}{72b^3} \\ & \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}}.\end{aligned}$$

Since $(bA)^{\frac{2}{b}} \geq \frac{1}{(b+1)^{\frac{2}{b}}} \geq \frac{1}{3} > \frac{1}{4}$ and $b \geq 2l$, we have

$$\begin{aligned}\eta_1 \geq {} & \frac{2l(l+b-3)(b+2l)}{12b^2}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\ & + \frac{l(b+2(l-1))(4(l-1)(2(l-1)-b)-3(2l-b)(l-b))}{72b^3} \\ & \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\ = {} & \frac{l\left[9b^3+(35l-26)b^2+(36l^2-90l)b+(4l^3-36l^2+48l-16)\right]}{72b^3} \\ & \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\ \geq {} & \frac{l\left[72l^3+(70l^2-52l)b+(36l^2-90l)b-36l^2\right]}{72b^3}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\ \geq {} & \frac{l\left[(72l^3-36l^2)+(140l-52l)b+(72l-90l)b\right]}{72b^3}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\ \geq {} & 0,\end{aligned}$$

which implies

$$f(\tau) \geq b(bA)^{\frac{b+2l}{b}} + \frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}} + \frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}.$$

Case 2: $2l-2 \leq b < 2l$. By using Taylor formula, we obtain the following inequalities for $t > 0$:

$$(1+t)^{\frac{2l}{b}} \geq 1 + \frac{2l}{b}t + \frac{2l(2l-b)}{2b^2}t^2 + \frac{2l(2l-b)(2l-2b)}{6b^3}t^3$$

and

$$\begin{aligned}(1+t)^{\frac{b+2(l-1)}{b}} \geq {} & 1 + \frac{2(l-1)+b}{b}t + \frac{(2(l-1)+b)(l-1)}{b^2}t^2 \\ & + \frac{(2(l-1)+b)(l-1)(2(l-1)-b)}{3b^3}t^3.\end{aligned}$$

Putting

$$t = \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}},$$

we have $b-2lt>\frac{l}{3}>0$,

$$\begin{aligned}
&\left(b-\frac{l(b+2(l-1))}{6(b+2l)}(bA)^{-\frac{2}{b}}\right)\left(1+\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{2l}{b}}\\
&=(b-2lt)(1+t)^{\frac{2l}{b}}\\
&\geq (b-2lt)\left[1+\frac{2l}{b}t+\frac{2l(2l-b)}{2b^2}t^2+\frac{2l(2l-b)(2l-2b)}{6b^3}t^3\right]\\
&=b-\frac{l(b+2l)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2-\frac{(2l-b)(8l^2+4lb)}{6b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\quad-\frac{4l^2(2l-b)(2l-2b)}{6b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4
\end{aligned}$$

and

$$\begin{aligned}
&\left(1+\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{b+2(l-1)}{b}}\\
&=(1+t)^{\frac{b+2(l-1)}{b}}\\
&\geq 1+\frac{2(l-1)+b}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)\\
&\quad+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\\
&\quad+\frac{(2(l-1)+b)(l-1)(2(l-1)-b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3.
\end{aligned}$$

Furthermore, we deduce by using the same method as the Case (1)

$$\begin{aligned}
f(\tau)&=(b+2l)\tau^{2l}bA-2l\tau^{b+2l}+\frac{l}{6}\tau^{b+2(l-1)}\\
&\geq (bA)^{\frac{b+2l}{b}}\left[b-\frac{l(b+2l)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\right.\\
&\quad-\frac{(2l-b)(8l^2+4lb)}{6b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\quad\left.-\frac{4l^2(2l-b)(2l-2b)}{6b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4\right]\\
&\quad+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}\left[1+\frac{2(l-1)+b}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)\right.\\
&\quad+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\\
&\quad\left.+\frac{(2(l-1)+b)(l-1)(2(l-1)-b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\right]\\
&=b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}+\eta_2,
\end{aligned}$$

where

$$\begin{aligned}
\eta_2 = & \frac{2l(l+b-3)(b+2l)}{3b^2}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-6}{b}} \\
& + \frac{l(2(l-1)+b)[(l-1)(2(l-1)-b)(b+2l)-l(2l-b)(2l-2b)]}{18b^3(b+2l)} \\
& \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
\geq & \frac{2l(l+b-3)(b+2l)}{9b^2}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
& + \frac{l(2(l-1)+b)(l-1)(2(l-1)-b)}{18b^3}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
= & \left[\frac{4bl(l+b-3)(b+2l)}{18b^3} + \frac{l(2(l-1)+b)(l-1)(2(l-1)-b)}{18b^3}\right] \\
& \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
\geq & \left[\frac{4bl(l+b-3)(b+2l)}{18b^3} + \frac{lb(b+2l)(2(l-1)-b)}{18b^3}\right] \\
& \times \left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
\geq & \frac{bl(b+2l)(6l+3b-14)}{18b^3}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3 (bA)^{\frac{b+2l-8}{b}} \\
\geq & \, 0
\end{aligned}$$

since $(bA)^{\frac{2}{b}} \geq \frac{1}{(b+1)^{\frac{2}{b}}} \geq \frac{1}{3}$. Therefore, we have

$$f(\tau) \geq b(bA)^{\frac{b+2l}{b}} + \frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}} + \frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}.$$

Case 3: $l \leq b < 2l-2$. By using the Taylor formula, one has for $t > 0$

$$(1+t)^{\frac{2l}{b}} \geq 1 + \frac{2l}{b}t + \frac{l(2l-b)}{b^2}t^2 + \frac{l(2l-b)(2l-2b)}{3b^3}t^3$$

and

$$(1+t)^{\frac{b+2(l-1)}{b}} \geq 1 + \frac{2(l-1)+b}{b}t + \frac{(2(l-1)+b)(l-1)}{b^2}t^2.$$

Putting

$$t = \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}} > 0,$$

one has $b - 2lt > 0$,

$$\begin{aligned}
&\left(b-\frac{l(b+2(l-1))}{6(b+2l)}(bA)^{-\frac{2}{b}}\right)\left(1+\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{2l}{b}}\\
&=(b-2lt)(1+t)^{\frac{2l}{b}}\\
&\geq (b-2lt)\left[1+\frac{2l}{b}t+\frac{l(2l-b)}{b^2}t^2+\frac{l(2l-b)(2l-2b)}{3b^3}t^3\right]\\
&=b-\frac{l(b+2l)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2-\frac{(2l-b)(4l^2+2lb)}{3b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\quad-\frac{4l^2(2l-b)(l-b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4
\end{aligned}$$

and

$$\begin{aligned}
\left(1+\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^{\frac{b+2(l-1)}{b}}&=(1+t)^{\frac{b+2(l-1)}{b}}\\
&\geq 1+\frac{2(l-1)+b}{b}\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\\
&\quad+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2.
\end{aligned}$$

By the same argument as the Case 2, we can deduce the following

$$\begin{aligned}
f(\tau)&=(b+2l)\tau^{2l}bA-2l\tau^{b+2l}+\frac{l}{6}\tau^{b+2(l-1)}\\
&\geq (bA)^{\frac{b+2l}{b}}\left[b-\frac{l(b+2l)}{b}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\right.\\
&\qquad-\frac{(2l-b)(4l^2+2lb)}{3b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^3\\
&\qquad\left.-\frac{4l^2(2l-b)(l-b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^4\right]\\
&\qquad+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}\left[1+\frac{2(l-1)+b}{b}\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right.\\
&\qquad\left.+\frac{(2(l-1)+b)(l-1)}{b^2}\left(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\right)^2\right]\\
&=b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}+\eta_3,
\end{aligned}$$

where

$$\begin{aligned}
\eta_3=&\frac{2l(l+b-3)(b+2l)}{3b^2}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^3(bA)^{\frac{b+2l-6}{b}}\\
&-\frac{4l^2(2l-b)(l-b)}{3b^3}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^4(bA)^{\frac{b+2l-8}{b}}\\
\geq&\, 0.
\end{aligned}$$

Therefore, we have

$$f(\tau) \geq b(bA)^{\frac{b+2l}{b}} + \frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}} + \frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}.$$

Case 4: $2 \leq b < l$. Since $2 \leq b < l$, there exists a positive integer k such that $2 \leq k-1 \leq \frac{2l}{b} < k$, then we have for $t > 0$ that

$$\begin{aligned}(1+t)^{\frac{2l}{b}} &\geq 1 + \frac{2l}{b}t + \frac{1}{2!}\frac{2l}{b}\left(\frac{2l}{b}-1\right)t^2 + \frac{1}{3!}\frac{2l}{b}\left(\frac{2l}{b}-1\right)\left(\frac{2l}{b}-2\right)t^3 \\ &\quad + \cdots + \frac{1}{(k+1)!}\frac{2l}{b}\left(\frac{2l}{b}-1\right)\cdots\left(\frac{2l}{b}-k\right)t^{k+1} \\ &= 1 + \sum_{p=0}^{k}\left\{\frac{1}{(p+1)!}\prod_{q=0}^{p}\left(\frac{2l}{b}-q\right)\right\}t^{p+1},\end{aligned}$$

$$\begin{aligned}(1+t)^{\frac{b+2l}{b}} &\leq 1 + \frac{b+2l}{b}t + \frac{1}{2!}\frac{b+2l}{b}\frac{2l}{b}t^2 + \frac{1}{3!}\frac{b+2l}{b}\frac{2l}{b}\left(\frac{2l}{b}-1\right)t^3 \\ &\quad + \cdots + \frac{1}{(k+1)!}\frac{b+2l}{b}\frac{2l}{b}\left(\frac{2l}{b}-1\right)\cdots\left(\frac{2l}{b}-(k-1)\right)t^{k+1} \\ &= 1 + \sum_{p=0}^{k}\left\{\frac{1}{(p+1)!}\prod_{q=0}^{p}\left(\frac{2l}{b}-q+1\right)\right\}t^{p+1},\end{aligned}$$

and

$$\begin{aligned}(1+t)^{\frac{b+2(l-1)}{b}} &\geq 1 + \frac{2(l-1)+b}{b}t + \frac{1}{2!}\frac{(2(l-1)+b)}{b}\frac{2(l-1)}{b}t^2 \\ &\quad + \frac{1}{3!}\frac{(2(l-1)+b)}{b}\frac{2(l-1)}{b}\left(\frac{2(l-1)}{b}-1\right)t^3 \\ &\quad + \cdots + \frac{1}{k!}\frac{(2(l-1)+b)}{b}\frac{2(l-1)}{b}\cdots\left(\frac{2(l-1)}{b}-(k-2)\right)t^k \\ &\quad - \left|\frac{1}{(k+1)!}\frac{(2(l-1)+b)}{b}\frac{2(l-1)}{b}\cdots\left(\frac{2(l-1)}{b}-(k-1)\right)\right|t^{k+1} \\ &= 1 + \sum_{p=0}^{k-1}\left\{\frac{1}{(p+1)!}\prod_{q=0}^{p}\left(\frac{2(l-1)}{b}-q+1\right)\right\}t^{p+1} \\ &\quad - \left|\frac{1}{(k+1)!}\prod_{q=0}^{k}\left(\frac{2(l-1)}{b}-q+1\right)\right|t^{k+1}.\end{aligned}$$

Putting $t = \frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}$ and $f(\tau) = (bA)^{\frac{b+2l}{b}}h(\tau)$, where

$$h(\tau) = (b+2l)(1+t)^{\frac{2l}{b}} - 2l(1+t)^{\frac{b+2l}{b}} + \frac{1}{6}(bA)^{-\frac{2}{b}}(1+t)^{\frac{b+2(l-1)}{b}},$$

then we have for $2 \leq b < l$,

$$
\begin{aligned}
h(\tau) \geq{}& (b+2l)\Bigg\{1+\sum_{p=0}^{k}\Bigg[\frac{1}{(p+1)!}\prod_{q=0}^{p}\Bigg(\frac{2l}{b}-q\Bigg)\Bigg]t^{p+1}\Bigg\} \\
&-2l\Bigg\{1+\sum_{p=0}^{k}\Bigg[\frac{1}{(p+1)!}\prod_{q=0}^{p}\Bigg(\frac{2l}{b}-q+1\Bigg)\Bigg]t^{p+1}\Bigg\} \\
&+\frac{l}{6}(bA)^{-\frac{2}{b}}\Bigg\{1+\sum_{p=0}^{k-1}\Bigg[\frac{1}{(p+1)!}\prod_{q=0}^{p}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg]t^{p+1} \\
&-\Bigg|\frac{1}{(k+1)!}\prod_{q=0}^{k}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg|t^{k+1}\Bigg\} \\
={}& b+\frac{l}{6}(bA)^{-\frac{2}{b}}+\sum_{p=1}^{k}\Bigg\{\frac{b+2l}{(p+1)!}\frac{2l}{b}\Bigg[\prod_{q=1}^{p}\Bigg(\frac{2l}{b}-q\Bigg)-\prod_{q=1}^{p}\Bigg(\frac{2l}{b}-q+1\Bigg)\Bigg]\Bigg\}t^{p+1} \\
&+\sum_{p=0}^{k-1}\Bigg\{\frac{l(bA)^{-\frac{2}{b}}}{6(p+1)!}\prod_{q=0}^{p}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg\}t^{p+1} \\
&-\Bigg|\frac{l(bA)^{-\frac{2}{b}}}{6(k+1)!}\prod_{q=0}^{k}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg|t^{k+1} \\
={}& b+\frac{l}{6}(bA)^{-\frac{2}{b}}-\sum_{p=1}^{k}\Bigg\{\frac{p2l(b+2l)}{b(p+1)!}\prod_{q=1}^{p-1}\Bigg(\frac{2l}{b}-q\Bigg)\Bigg\}t^{p+1} \\
&+\sum_{p=1}^{k}\Bigg\{\frac{l(bA)^{-\frac{2}{b}}}{6p!}\prod_{q=0}^{p-1}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg\}t^{p} \\
&-\Bigg|\frac{l(bA)^{-\frac{2}{b}}}{6(k+1)!}\prod_{q=0}^{k}\Bigg(\frac{2(l-1)}{b}-q+1\Bigg)\Bigg|t^{k+1} \\
={}& b+\frac{l}{6}(bA)^{-\frac{2}{b}}-\sum_{p=1}^{k}\Bigg\{\frac{p}{b^{p}(p+1)!}\prod_{q=0}^{p}(2l-(q-1)b)\Bigg\}\Bigg(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\Bigg)^{p+1} \\
&+\sum_{p=1}^{k}\Bigg\{\frac{l(bA)^{-\frac{2}{b}}}{6b^{p}p!}\prod_{q=0}^{p-1}(2(l-1)-(q-1)b)\Bigg\}\Bigg(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\Bigg)^{p} \\
&-\Bigg|\frac{l(bA)^{-\frac{2}{b}}}{6b^{k+1}(k+1)!}\prod_{q=0}^{k}(2(l-1)-(q-1)b)\Bigg|\Bigg(\frac{b+2(l-1)}{12(b+2l)}(bA)^{-\frac{2}{b}}\Bigg)^{k+1}.
\end{aligned}
$$

Furthermore,

$$
\begin{aligned}
f(\tau) \geq{}& b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^{2}}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}} \\
&-\sum_{p=2}^{k}\Bigg\{\frac{p}{b^{p}(p+1)!}\prod_{q=0}^{p}(2l-(q-1)b)\Bigg\}\Bigg(\frac{b+2(l-1)}{12(b+2l)}\Bigg)^{p+1}(bA)^{\frac{b+2l-2p-2}{b}}
\end{aligned}
$$

$$\begin{aligned}
&+\sum_{p=2}^{k}\left\{\frac{l}{6b^pp!}\prod_{q=0}^{p-1}(2(l-1)-(q-1)b)\right\}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^p(bA)^{\frac{b+2l-2p-2}{b}}\\
&-\left|\frac{l}{6b^{k+1}(k+1)!}\prod_{q=0}^{k}(2(l-1)-(q-1)b)\right|\left(\frac{b+2(l-1)}{12(b+2l)}\right)^{k+1}(bA)^{\frac{b+2l-2k-4}{b}}\\
=\;&b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}+\eta_4,
\end{aligned}$$

where

$$\begin{aligned}
\eta_4=&\sum_{p=2}^{k}\left\{\frac{(b+2(l-1))2l}{12b^pp!}\left[\prod_{q=1}^{p-1}(2(l-1)-(q-1)b)-\frac{p}{p+1}\prod_{q=1}^{p-1}(2l-qb)\right]\right\}\\
&\times\left(\frac{b+2(l-1)}{12(b+2l)}\right)^p(bA)^{\frac{b+2l-2p-2}{b}}\\
&-\left|\frac{l}{6b^{k+1}(k+1)!}\prod_{q=0}^{k}(2(l-1)-(q-1)b)\right|\left(\frac{b+2(l-1)}{12(b+2l)}\right)^{k+1}(bA)^{\frac{b+2l-2k-4}{b}}.
\end{aligned}$$

From $k-2\le\frac{2(l-1)}{b}<k$, we have

$$\frac{k-2-i}{k+1-i}\le\frac{\frac{2(l-1)}{b}-i}{k+1-i}<\frac{k-i}{k+1-i}. \tag{2.7}$$

Then, it follows from (2.7) that

$$\left|\frac{\frac{2(l-1)}{b}-i}{k+1-i}\right|\le 1,\quad \text{for } i=0,1,2,\cdots,k-1.$$

Note that

$$2(l-1)-(q-1)b\ge 2l-qb\ge 0,\quad \text{for } p=2,3,\cdots,k,$$

one has

$$\begin{aligned}
&\prod_{q=1}^{p-1}(2(l-1)-(q-1)b)-\frac{p}{p+1}\prod_{q=1}^{p-1}(2l-qb)\\
\ge&\prod_{q=1}^{p-1}(2(l-1)-(q-1)b)-\prod_{q=1}^{p-1}(2l-qb)\ge 0.
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
&\sum_{p=2}^{k}\left\{\frac{(b+2(l-1))2l}{12b^pp!}\left[\prod_{q=1}^{p-1}(2(l-1)-(q-1)b)-\frac{p}{p+1}\prod_{q=1}^{p-1}(2l-qb)\right]\right\}\\
&\quad\times\left(\frac{b+2(l-1)}{12(b+2l)}\right)^p(bA)^{\frac{b+2l-2p-2}{b}}\\
\ge&\frac{(b+2(l-1))2l}{24b^2}\left[2(l-1)-\frac{2(2l-b)}{3}\right]\left(\frac{b+2(l-1)}{12(b+2l)}\right)^2(bA)^{\frac{b+2l-6}{b}}.
\end{aligned}$$

From

$$(bA)^{\frac{2}{b}}\ge\frac{1}{(b+1)^{\frac{2}{b}}}\ge\frac{1}{3},$$

we have

$$\begin{aligned}
\eta_4 \geq\ & \frac{(b+2(l-1))2l}{24b^2}\left[2(l-1)-\frac{2(2l-b)}{3}\right]\left(\frac{b+2(l-1)}{12(b+2l)}\right)^2(bA)^{\frac{b+2l-6}{b}} \\
& -\left|\frac{l}{6b^{k+1}(k+1)!}\prod_{q=0}^{k}(2(l-1)-(q-1)b)\right|\left(\frac{b+2(l-1)}{12(b+2l)}\right)^{k+1}(bA)^{\frac{b+2l-2k-4}{b}} \\
=\ & \frac{l(b+2(l-1))}{12b^2}\Bigg\{\left[2(l-1)-\frac{2(2l-b)}{3}\right] \\
& -2b\left|\frac{1}{b^k(k+1)!}\prod_{q=1}^{k}(2(l-1)-(q-1)b)\right|\left(\frac{b+2(l-1)}{12(b+2l)}\right)^{k-1}(bA)^{\frac{-2k+2}{b}}\Bigg\} \\
& \times\left(\frac{b+2(l-1)}{12(b+2l)}\right)^2(bA)^{\frac{b+2l-6}{b}} \\
\geq\ & \frac{l(b+2(l-1))}{12b^2}\left\{\frac{2l+2b-6}{3}-2b\left|\prod_{q=0}^{k-1}\left(\frac{\frac{2(l-1)}{b}-q}{k+1-q}\right)\right|\left(\frac{1}{4}\right)^{k-1}\right\} \\
& \times\left(\frac{b+2(l-1)}{12(b+2l)}\right)^2(bA)^{\frac{b+2l-6}{b}} \\
\geq\ & \frac{l(b+2(l-1))}{12b^2}\left\{\frac{2l+2b-6}{3}-\frac{b}{8}\right\}\left(\frac{b+2(l-1)}{12(b+2l)}\right)^2(bA)^{\frac{b+2l-6}{b}} \\
\geq\ & 0,
\end{aligned}$$

which implies that

$$f(\tau)\geq b(bA)^{\frac{b+2l}{b}}+\frac{l}{6}(bA)^{\frac{b+2(l-1)}{b}}+\frac{l(b+2(l-1))^2}{144b(b+2l)}(bA)^{\frac{b+2l-4}{b}}.$$

This completes the proof of Lemma 2.1.

□

3 Proof of Theorem 1.1 and Remark 1.2

Proof of Theorem 1.1. We will use the same notations as those of [3]. In this section, we assume that $b=n$. Let $\widehat{\varphi}_j(z)$ be the Fourier transform of the trial function $\varphi_j(x)$,

$$\varphi_j(x)=\begin{cases}u_j(x), & x\in\Omega,\\ 0, & x\in\mathbb{R}^n\setminus\Omega,\end{cases}$$

where $u_j(x)$ is an orthonormal eigenfunction corresponding to the eigenvalue λ_j, $f(z):=\sum_{j=1}^{k}|\widehat{\varphi}_j(z)|^2$, and f^* be the symmetric decreasing rearrangement of f. And then, we can obtain from Lemma 2.1 that

$$\begin{aligned}\sum_{j=1}^{k}\lambda_j &\geq nB_n\int_0^\infty s^{n+2l-1}\phi(s)ds\\ &\geq \frac{nB_n\left(\frac{k}{B_n}\right)^{\frac{n+2l}{n}}}{n+2l}\phi(0)^{-\frac{2l}{n}}+\frac{lB_n\left(\frac{k}{B_n}\right)^{\frac{n+2(l-1)}{n}}}{6(n+2l)\mu^2}\phi(0)^{\frac{2n-2l+2}{n}}\\ &\quad+\frac{l(n+2(l-1))^2B_n\left(\frac{k}{B_n}\right)^{\frac{n+2l-4}{n}}}{144n(n+2l)^2\mu^4}\phi(0)^{\frac{4n-2l+4}{n}},\end{aligned}\tag{3.1}$$

where $\phi:[0,\ +\infty)\to[0,\ (2\pi)^{-n}V(\Omega)]$ is a non-increasing function of $|x|$ and $\phi(x)$ is defined by $\phi(|x|):=f^*(x)$. Now defining a function $\xi(t)$ as follows:

$$\begin{aligned}\xi(t)&=\frac{nB_n}{n+2l}\left(\frac{k}{B_n}\right)^{\frac{n+2l}{n}}t^{-\frac{2l}{n}}+\frac{lB_n}{6(n+2l)\mu^2}\left(\frac{k}{B_n}\right)^{\frac{n+2(l-1)}{n}}t^{\frac{2n-2l+2}{n}}\\ &\quad+\frac{l(n+2(l-1))^2B_n}{144n(n+2l)^2\mu^4}\left(\frac{k}{B_n}\right)^{\frac{n+2l-4}{n}}t^{\frac{4n-2l+4}{n}}.\end{aligned}\tag{3.2}$$

Here we assume that $l\leq n+1$. The other cases (i.e., $n+1<l<2(n+1)$, $l\geq 2(n+1)$) can be discussed by using of the similar method. After differentiating (3.2) with respect to the variable t, we derive

$$\begin{aligned}\xi'(t)&=\frac{B_n}{n+2l}\left(\frac{k}{B_n}\right)^{\frac{n+2l}{n}}t^{-\frac{2l}{n}-1}\Bigg[-2l+\frac{l(2n-2l+2)}{6n\mu^2}\left(\frac{k}{B_n}\right)^{-\frac{2}{n}}t^{\frac{2n+2}{n}}\\ &\quad+\frac{l(4n-2l+4)(n+2(l-1))^2}{144n^2(n+2l)\mu^4}\left(\frac{k}{B_n}\right)^{-\frac{4}{n}}t^{\frac{4n+4}{n}}\Bigg].\end{aligned}\tag{3.3}$$

Putting $\zeta(t)=\xi'(t)\frac{n+2l}{B_n}(\frac{k}{B_n})^{-\frac{n+2l}{n}}t^{\frac{2l}{n}+1}$ and noticing that $\mu\geq(2\pi)^{-n}B_n^{-\frac{1}{n}}V(\Omega)^{\frac{n+1}{n}}$, we can deduce from (3.3)

$$\begin{aligned}\zeta(t)&=-2l+\frac{l(2n-2l+2)}{6n\mu^2}\left(\frac{k}{B_n}\right)^{-\frac{2}{n}}t^{\frac{2n+2}{n}}\\ &\quad+\frac{l(4n-2l+4)(n+2(l-1))^2}{144n^2(n+2l)\mu^4}\left(\frac{k}{B_n}\right)^{-\frac{4}{n}}t^{\frac{4n+4}{n}}\\ &\leq -2l+\frac{l(2n-2l+2)}{6n(2\pi)^{-2n}B_n^{-\frac{2}{n}}Vol(\Omega)^{\frac{2(n+1)}{n}}}\left(\frac{k}{B_n}\right)^{-\frac{2}{n}}t^{\frac{2n+2}{n}}\\ &\quad+\frac{l(4n-2l+4)(n+2(l-1))^2}{144n^2(n+2l)(2\pi)^{-4n}B_n^{-\frac{4}{n}}Vol(\Omega)^{\frac{4(n+1)}{n}}}\left(\frac{k}{B_n}\right)^{-\frac{4}{n}}t^{\frac{4n+4}{n}}.\end{aligned}\tag{3.4}$$

Since the right hand side of (3.4) is an increasing function of t, if the right hand side of (3.4) is not larger than 0 at $t = (2\pi)^{-n}V(\Omega)$, that is

$$\begin{aligned}\zeta(t) \le& -2l + \frac{l(2n-2l+2)}{6n}k^{-\frac{2}{n}}\frac{B_n^{\frac{4}{n}}}{(2\pi)^2} \\ &+ \frac{l(4n-2l+4)(n+2(l-1))^2}{144n^2(n+2l)}k^{-\frac{4}{n}}\frac{B_n^{\frac{8}{n}}}{(2\pi)^4} \\ \le& \, 0,\end{aligned} \tag{3.5}$$

we can claim from (3.5) that $\xi'(t) \le 0$ on $(0, (2\pi)^{-n}V(\Omega)]$. If $\xi'(t) \le 0$, then $\xi(t)$ is a decreasing function on $(0, (2\pi)^{-n}V(\Omega)]$. In fact, by a direct calculation, we can obtain

$$\zeta(t) \le -2l + \frac{l(2n-2l+2)}{6n} + \frac{l(4n-2l+4)(n+2(l-1))^2}{144n^2(n+2l)} \le 0 \tag{3.6}$$

since $\frac{B_n^{\frac{4}{n}}}{(2\pi)^2} < 1$.

On the other hand, since $0 < \phi(0) \le (2\pi)^{-n}V(\Omega)$ and right hand side of the formula (3.1) is $\xi(\phi(0))$, which is a decreasing function of $\phi(0)$ on $(0, (2\pi)^{-n}V(\Omega)]$, then we can replace $\phi(0)$ by $(2\pi)^{-n}V(\Omega)$ in (3.1) which gives the inequality as follows:

$$\begin{aligned}\frac{1}{k}\sum_{j=1}^{k}\lambda_j \ge& \frac{n}{n+2l}\frac{(2\pi)^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}} \\ &+ \frac{l}{24(n+2l)}\frac{(2\pi)^{2(l-1)}}{(B_nV(\Omega))^{\frac{2(l-1)}{n}}}\frac{V(\Omega)}{I(\Omega)}k^{\frac{2(l-1)}{n}} \\ &+ \frac{l(n+2(l-1))^2}{2304n(n+2l)^2}\frac{(2\pi)^{2(l-2)}}{(B_nV(\Omega))^{\frac{2(l-2)}{n}}}\left(\frac{V(\Omega)}{I(\Omega)}\right)^2 k^{\frac{2(l-2)}{n}}.\end{aligned}$$

This completes the proof of Theorem 1.1.

□

Next we will prove that the inequality (1.17) is sharper than the inequality (1.16).

Proof of Remark 1.2: Under the same assumption with Lemma 2.1, let $b = n$ and $A = \frac{k}{nB_n}$, we obtain from $\mu \ge (2\pi)^{-n}B_n^{-\frac{1}{n}}V(\Omega)^{\frac{n+1}{n}}$ that

$$\begin{aligned}\frac{(bA)^{-\frac{2}{b}}\psi(0)^{2+\frac{2}{b}}}{\mu^2} \le& \frac{(2\pi)^{-2n-2}V(\Omega)^{2+\frac{2}{n}}}{(2\pi)^{-2n}B_n^{-\frac{2}{n}}V(\Omega)^{\frac{2(n+1)}{n}}}(\frac{k}{B_n})^{-\frac{2}{n}} \\ =& \frac{(2\pi)^{-2}}{(B_n)^{-\frac{4}{n}}}k^{-\frac{2}{n}} < \frac{(2\pi)^{-2}}{(B_n)^{-\frac{4}{n}}} < 1,\end{aligned}$$

then we have

$$\begin{aligned}&\frac{1}{b+2l}\sum_{p=2}^{l}\frac{(l+1-p)}{(6)^p b\cdots(b+2p-2)\mu^{2p}}(bA)^{\frac{b+2(l-p)}{b}}\psi(0)^{\frac{2pb-2(l-p)}{b}} \\ <& \frac{1}{b+2l}\sum_{p=2}^{l}\frac{(l+1-p)}{(6)^p b\cdots(b+2p-2)\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}\end{aligned}$$

$$\begin{aligned}
&< \frac{l-1}{36(b+2l)b(b+2)\mu^4}\sum_{p=0}^{\infty}\frac{1}{6^p(b+2)^p}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}\\
&= \frac{l-1}{6b(b+2l)(6(b+2)-1)\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}.
\end{aligned} \tag{3.7}$$

By a direct calculation, we derive

$$l(6(b+2)-1)(b+2(l-1))^2 > 24b(b+2l)(l-1) > 0,$$

in fact,

$$\begin{aligned}
&l(6(b+2)-1)(b+2(l-1))^2 - 24b(b+2l)(l-1)\\
&= 4b(l-1)[6b(l-1)-l] + l(6b+11)[b^2+4(l-1)^2]\\
&> 24b^2(l-1)^2 + 4bl(l-1)[6(l-1)-1] > 0,
\end{aligned}$$

that is,

$$\frac{24b(b+2l)(l-1)}{l(6(b+2)-1)(b+2(l-1))^2} < 1. \tag{3.8}$$

Therefore, we get from (3.7) and (3.8) that

$$\begin{aligned}
&\frac{1}{b+2l}\sum_{p=2}^{l}\frac{(l+1-p)}{(6)^p b\cdots(b+2p-2)\mu^{2p}}(bA)^{\frac{b+2(l-p)}{b}}\psi(0)^{\frac{2pb-2(l-p)}{b}}\\
&< \frac{l-1}{6b(b+2l)(6(b+2)-1)\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}\\
&= \frac{24b(b+2l)(l-1)}{l(6(b+2)-1)(b+2(l-1))^2}\cdot\frac{l(b+2(l-1))^2}{144b^2(b+2l)^2\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}\\
&< \frac{l(b+2(l-1))^2}{144b^2(b+2l)^2\mu^4}(bA)^{\frac{b+2l-4}{b}}\psi(0)^{\frac{4b-2l+4}{b}}.
\end{aligned} \tag{3.9}$$

Taking

$$b=n,\quad A=\frac{k}{nB_n},\quad \psi(0)=(2\pi)^{-n}V(\Omega),\quad \mu=2(2\pi)^{-n}\sqrt{V(\Omega)I(\Omega)}, \tag{3.10}$$

and substituting (3.10) into (3.9), one has

$$\begin{aligned}
&\frac{n}{n+2l}\frac{(2\pi)^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}} + \frac{l}{24(n+2l)}\frac{(2\pi)^{2(l-1)}}{(B_nV(\Omega))^{\frac{2(l-1)}{n}}}\frac{V(\Omega)}{I(\Omega)}k^{\frac{2(l-1)}{n}}\\
&+\frac{l(n+2(l-1))^2}{2304n(n+2l)^2}\frac{(2\pi)^{2(l-2)}}{(B_nV(\Omega))^{\frac{2(l-2)}{n}}}\left(\frac{V(\Omega)}{I(\Omega)}\right)^2 k^{\frac{2(l-2)}{n}}\\
>&\frac{n}{n+2l}\frac{(2\pi)^{2l}}{(B_nV(\Omega))^{\frac{2l}{n}}}k^{\frac{2l}{n}} + \frac{n}{(n+2l)}\\
&\times\sum_{p=1}^{l}\frac{l+1-p}{(24)^p n\cdots(n+2p-2)}\frac{(2\pi)^{2(l-p)}}{(B_nV(\Omega))^{\frac{2(l-p)}{n}}}\left(\frac{V(\Omega)}{I(\Omega)}\right)^p k^{\frac{2(l-p)}{n}}.
\end{aligned} \tag{3.11}$$

This completes the proof of Remark 1.2.

□

Acknowledgments The authors wish to express their gratitude to Prof. Q. -M. Cheng for continuous encouragement and enthusiastic help.

References

[1] S. Agmon, On kernels, eigenvalues and eigenfunctions of operators related to elliptic problems, Comm. Pure Appl. Math. 18 (1965), 627-663.

[2] F. A. Berezin, Covariant and contravariant symbols of operators, Izv. Akad. Nauk SSSR Ser. Mat. 36 (1972), 1134-1167.

[3] Q. -M. Cheng, X. Qi and G. Wei, A lower bound for eigenvalues of the poly-Laplacian with arbitrary order, http://arxiv.org/abs/1012.3006v1.

[4] Q. -M. Cheng and G. Wei, A lower bound for eigenvalues of a clamped plate problem, Calc. Var. Partial Differential Equations 42 (2011), 579-590.

[5] Q. -M. Cheng and G. Wei, Upper and lower bounds of the clamped plate problem, preprint, (2011).

[6] A. Laptev, Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces, J. Funct. Anal. 151 (1997), 531-545.

[7] H. A. Levine and M. H. Protter, Unrestricted lower bounds for eigenvalues for classes of elliptic equations and systems of equations with applications to problems in elasticity, Math. Methods Appl. Sci. 7 (1985), no. 2, 210-222.

[8] P. Li and S. T. Yau, On the Schrödinger equations and the eigenvalue problem, Comm. Math. Phys. 88 (1983), 309-318.

[9] E. Lieb, The number of bound states of one-body Schrödinger operators and the Weyl problem, Proc. Sym. Pure Math. 36 (1980), 241-252.

[10] A. D. Melas, A lower bound for sums of eigenvalues of the Laplacian, Proc. Amer. Math. Soc. 131 (2003), 631-636.

[11] A. Pleijel, On the eigenvalues and eigenfunctions of elastic plates, Comm. Pure Appl. Math., 3 (1950), 1-10.

[12] G. Pólya, On the eigenvalues of vibrating membranes, Proc. London Math. Soc., 11 (1961), 419-433.

[13] J. Jost, X. Li-Jost, Q. Wang, and C. Xia, Universal bounds for eigenvalues of the polyharmonic operators, Trans. Amer. Math. Soc., 363(2011), 1821-1854.

Guoxin Wei, School of Mathematical Sciences, South China Normal University,
510631, Guangzhou, China, weigx03@mails.tsinghua.edu.cn
Lingzhong Zeng, Department of Mathematics, Graduate School of Science and
Engineering, Saga University, Saga 840-8502, Japan, lingzhongzeng@yeah.net